

Galliprant[®]
(grapiprant tablets)

HENRY SCHEIN[®]
ANIMAL HEALTH
(855) SCHEIN1 (724-3461)
www.henryscheinvet.com
(888) 329-3861 fax
Order through your distributor!

ONE GIANT LEAP
FOR MAN'S BEST FRIEND



INTRODUCING GALLIPRANT[®] (grapiprant tablets)

TREAT OSTEOARTHRITIS (OA) WHEN DOGS NEED IT

One key to your canine patients' health is helping them stay active. Start targeted pain relief from the earliest diagnosed stages of canine OA with Galliprant. Give these dogs the relief they need and help keep them doing the things they love.

TREAT WITH GALLIPRANT FROM THE EARLIEST DIAGNOSED STAGES OF CANINE OA

- Galliprant is a piperidine class drug for a new class of canine OA therapy
- Mode of action targets canine OA pain and inflammation while reducing the impact on GI, kidney and liver homeostasis^{1,2}
- Proven safe
- Most dogs can be dosed with a whole or half tablet*

*Dosage should be calculated in half-tablet increments. Dogs weighing less than 8 lbs (3.6 kg) cannot be accurately dosed.

GALLIPRANT PRICING

	Size (mg)	Tablets per bottle	Price per bottle
	20 mg	30	\$24.00
	60 mg	30	\$48.30
	100 mg	30	\$72.60
	20 mg	90	\$72.00
	60 mg	90	\$144.90
	100 mg	90	\$217.80

CONTACT YOUR ELANCO, ARATANA OR DISTRIBUTOR REPRESENTATIVE TO INCORPORATE GALLIPRANT INTO YOUR CANINE OA PAIN PROTOCOL TODAY

To learn more, please visit galliprantfordogs.com/vet

Galliprant is indicated for the control of pain and inflammation associated with osteoarthritis in dogs.

IMPORTANT SAFETY INFORMATION

Not for use in humans. For use in dogs only. Keep this and all medications out of reach of children and pets. Store out of reach of dogs and other pets in a secured location in order to prevent accidental ingestion or overdose. Do not use in dogs that have a hypersensitivity to grapiprant. If Galliprant is used long term, appropriate monitoring is recommended. Concomitant use of Galliprant with other anti-inflammatory drugs, such as COX-inhibiting NSAIDs or corticosteroids, should be avoided. Concurrent use with other anti-inflammatory drugs or protein-bound drugs has not been studied. The safe use of Galliprant has not been evaluated in dogs younger than 9 months of age and less than 8 lbs (3.6 kg), dogs used for breeding, pregnant or lactating dogs, or dogs with cardiac disease. The most common adverse reactions were vomiting, diarrhea, decreased appetite, and lethargy. Please see enclosed product label or call 1-888-545-5973 for full prescribing information.

1. Kirkby Shaw, K., Rausch-Derra, L., and Rhodes, L. 2016. "Grapiprant: an EP4 prostaglandin receptor antagonist and novel therapy for pain and inflammation." *Vet. Med. Sci.* 2: 3-9.
2. Rausch-Derra, L., Huebner, M., and Rhodes, L. 2015. "Evaluation of the safety of long-term, daily oral administration of grapiprant, a novel drug for treatment of osteoarthritis pain and inflammation, in healthy dogs." *Am. J. Vet. Res.* 76.10: 853-859.

ARATANA
THERAPEUTICS

Elanco

